



Accuracy of Fine Needle Aspiration Biopsy (F.N.A.B) in Diagnosis of Breast Lump

¹ Sarab K. Abedalrahman, ² Najim Abid Issa Al-Khalidy, ³ Ali S. Al-Hashimi
⁴ Jawad K. Al-Diwan



ABSTRACT

Background: Breast cancer is the first one among Iraqi females. Most of them present later for diagnosis. Early detection center in tertiary hospital practice uses FNAB for early diagnosis. Publications on accuracy of this detection are scarce.

Objective: To test the accuracy of FNAB in breast lump diagnosis.

Methods: Diagnostic test accuracy study, on 204 women with breast lump, attending the oncology department in 2017.

Results: Fine-needle aspiration biopsy diagnosis of histologically malignant cases were, malignant in 89 (87.3%), suspicious of malignancy in 5 (4.9%), and benign in 4 (3.9%). Complete sensitivity was 87.3%, and specificity was 100%, with 12.7% false negative results and no false positive cases. The accuracy was 94%.

Conclusion: The study confirmed the high figures of accuracy of FNAB.

¹Corresponding Author to:

Al-Alwiyaa Maternity Teaching Hospital, women health center
e- mail: sarab.abedalrahman@yahoo.com

² Dept. of Family and Community Medicine, College of Medicine, University of Baghdad

Key words: breast cancer, fine needle aspiration, sensitivity, specificity.

Received at 15/10/2018

Accepted at 28/1/2019

INTRODUCTION

Breast cancer is commonest cancer among women, comprising about 23% of the newly diagnosed cancers among females, and 6 % of all deaths worldwide ^(1,2).

In Iraq, female incidence rate was 23.01 per 100,000, accounting about 34% of all reported cancer among females ⁽³⁾. Breast cancer survival rates tend to be poor in developing countries, most likely because of a combination of late diagnosis and limited access to timely and appropriate treatment ⁽⁴⁾. Iraqi patients presented at stage II, III, IV. ⁽⁵⁾ It was explained by low index of the suspicion and use of traditional methods ⁽⁶⁾.

Breast fine-needle aspiration biopsy (FNAB) of the breast is a minimally invasive diagnostic method, often obviating an open biopsy. ⁶ Ultrasound guided FNAB has been widely used for non-palpable breast lumps. ^(7,8) Although core biopsy is preferred to FNAB in most developed countries, its procedure is more expensive and time consuming as compared to FNAB ⁽⁹⁾.

The clinical use of FNAB has been questioned because of the variation in results ⁽¹⁰⁾.

A population based breast cancer screening program is not established in Iraq and only early detection centers, and clinics, were established at tertiary hospitals ¹².

Publishing data on the accuracy of FNAB in Iraq is scarce. The situation of screening was the impetus to carry out this study, it is objectives to evaluate the sensitivity, and specificity of FNAB.

Materials and methods

A total of 204 female patients were included in the study. They were recruited from women health centre in Al-Elwiyaa Maternity Teaching Hospital for the period Jan. 2017 – Nov. 2017. All selected women were subjected for FNAB. Histological examination was done after excisional biopsy or mastectomy.

FNAB techniques was carried out according to the suggested Iraqi guidelines ⁽¹³⁾.

Accuracy of FNAB was done by sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV). The sensitivity and specificity is the proportion of individuals that correctly diagnosed as diseased and non –diseased, respectively. The performance of the FNAB was tested by the PPV, NPV, Sensitivity, and Specificity ⁽¹⁴⁾.

RESULTS

Out of the total, 102 (50%) malignant cases, were diagnosed. The age of malignant and benign cases was 50.7(±12.2), and 33.7(±12.4), respectively. There was a significant difference in age between

malignant and benign cases ($t=9.834$, $df=202$, $P=0.001$).

Eighty-nine (87.3%), 5(4.9%), 4(3.9%), were diagnosed as malignant, suspicious of malignancy and benign, respectively (table

1). Out of those benign cases, there were 6 (5.9%), 11(10.8%), and 85(83.3%) as suspicious of malignancy, Atypical probably benign and benign, respectively.

Table 1: The histological (reference) and FNAB (index) tests result presentation

FNA	Histology result		Total
	Malignant	Benign	
Malignant	89 (87.3%)	0 (0%)	88 (43.1%)
Suspicious of malignancy	5 (4.9%)	6 (5.9%)	11(5.4%)
Atypical probably benign	4 (3.9%)	11(10.8%)	16(7.8%)
Total	120	102	204

$X^2=166.1$, $df=3$, $P= 0.00$

Sensitivity, specificity, PPV and NPV were 87.3%, 100%, 100% and 88.7%, respectively, as shown in table 2.

Table 2: The complete sensitivity and specificity of FNA

FNAB	Histological examination		Total
	Positive	Negative	
Positive	89	0	89
Negative	13	102	115
Total	102	102	204

Sensitivity = 87.3%, Specificity = 100%,
 PPV = 100%, NPV = 88.7%,
 Accuracy=94% FP = 0%, FN=12.7%,

DISCUSSION

For a good therapeutic management, there is a need for rapid, inexpensive, and non-invasive test. FNAB is the test of choice for this purpose because of the finer needle size and is easier/safer in certain lesions, such as very small lesions, lesions just under the skin or very close to the chest wall compared with true cut biopsy. In addition, FNAB maintains tactile sensitivity, allows multidirectional passes allowing a broader sampling of the lesion and immediate reporting where necessary. However, FNAB is less reliable at differentiating invasive cancer from DCIS, may be limited in some cases in the assessment of tumor grade and prognostic.⁽¹⁵⁾ The age of cases with malignant lesions was 50.7 ± 12.2 year. It is consistent with that reported in Iraq (49.4 ± 11.66) years¹⁶.

This study showed that sensitivity of FNAB was 87.3%. It is lower than that reported in literature (92.7%) in Meta-analysis study.¹⁷ In the recent years automated core needle biopsy, and various suction assisted devices of increasing core diameter, with mounting numbers of tissue pieces collected from each lesion, have replaced FNAB as the 1st line of diagnosis. The accuracy of FNAB depends on the experience of cytopathologist and the physician performing the procedure. High figures of accuracy reported in the world e.g. sensitivity (92%)⁽¹⁸⁾, (99.4%)⁽¹⁹⁾ and specificity (97%)¹⁸(100%)¹⁹.

The observed figure of sensitivity (87.3%) is within the reported sensitivity range in United Kingdom (84%-93%). The calculation of observed figure was complete one i.e. considering all abnormal results in one category. However, result of FNAB in UK used distinguished classification and results. The finding was outcome of the screening established in 1987⁽²⁰⁾. In Iraq, there are an early detection services rather than screening program¹².

In early years of breast screening program, FNAB was used, then Core biopsy (CB) was introduced and recently, wide bore vacuum biopsy. CB has been improved and considered the standard, completely in UK.²¹ Unfortunately in Iraq research still is digging in FNAB. The positive and negative predictive values of FNAB were 100%, and 88.7% respectively. Similar finding were reported in literature^(22, 23, 24). The FNAB is a good test for pre-operative decision, but true

cut always provides a better histological diagnosis and is more accurate and more trusted.

The suspicious diagnosis was found among 17 (13.2%) of the study subjects. This was lower than reported in Pakistan, (20%)²² and higher than that reported in Nigeria (12.4%).¹⁹ These variation might reflect the difficulties in the developing world, facing diagnosis of malignancy.

FNAB findings influenced by the maneuver (number of times inside the lump). It was practiced as 10 times maneuver to get harvests cells for examination^{25, 26}.

CONCLUSION

The study confirmed the high figures of accuracy of FNAB.

REFERENCES

1. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics 2002. *CA Cancer J Clin* 2005; 55(2):74–108.
2. Parkin DM, Fernandez LM. Use of statistics to assess the global burden of breast cancer. *Breast J*. 2006; 12(1 Suppl.):S70–80.
3. Iraqi Cancer Board, Ministry of Health (Iraq). Iraq Cancer Registry 2011. Baghdad, Iraq: Ministry of Health (Iraq); 2014.
4. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin*. 2011; 61:69-90. <http://dx.doi.org/10.3322/caac.2010>
5. Alwan NAS, Tawfeeq FN, Muallah MH, Sattar SA, Saleh WA. The Stage of Breast Cancer at the Time of Diagnosis: Correlation with the Clinicopathological Findings among Iraqi Patients. *J Neoplasm* 2017. 2:1-10
6. Gupta RK, Naran S, Buchanan A, Fauck R, Simpson J: Fine-needle aspiration cytology of breast: its impact on surgical practice with an emphasis on the diagnosis of breast abnormalities in young women. *Diagn Cytopathol* 1988; 4:206-209.
7. Evans WP, Cade SH. Needle localization and fine-needle aspiration biopsy of non-palpable breast lesions with use of standard and stereotactic equipment. *Radiology* 1989;173: 53-6.
8. Sterrett G, Oliver D, Frayne J, Ingram D, Sheiner H. Stereotactic fine needle aspiration biopsy (SFNB) of breast: preliminary results in Perth with the TRC mammo-test machine. Cytological aspects. *Pathology* 1991;23:302-10. <http://dx.doi.org/10.1111/j.1445-2197.1992.tb07226.x>
9. Lieu D. Value of cytopathologist-performed ultrasound-guided fine-needle aspiration as a screening test for ultrasound-guided core-needle biopsy in non-palpable breast masses. *Diagn Cytopathol*. 2009; 37: 262-9. [<http://dx.doi.org/10.1002/dc.20984>]
10. Jackson VP, Reynolds HE: Stereotactic needle-core biopsy and fine-needle aspiration cytologic evaluation of non-palpable breast lesions. *Radiology* 1991; 181:633-4.
11. IMOH. Iraq cancer committee. Early cancer detection. 9th oct. 2018. <https://www.moh.gov.iq>
12. Al Alwan Nada A. S. Establishing national guidelines for early detection of breast cancer in Iraq: Clinical Implications and Perspectives. *International Journal of Advanced Research* 2015;3(12): 539–55.
13. Eusebi P. Diagnostic Accuracy Measures. *Cerebrovasc Dis* 2013; 36: 267–72.
14. He Q, Fan X, Yuan T, Kong L, Du X, Zhuang D, Fan Z. Eleven years of experience reveals that fine-needle aspiration cytology is still a useful method for preoperative diagnosis of breast carcinoma. *Breast*. 2007; 16(3):303-6, <http://dx.doi.org/10.1016/j.breast.2006.12.006>.
15. Molah SA, Ali HH, Mohammed S A, Rahim F H. The incidence, age at diagnosis of breast cancer in the Iraqi Kurdish population and comparison to some other countries of Middle-East and West. *Int Jof Surg*. 2015; 13: 71-5.
16. Yu YH, Wei W, Liu JL. Diagnostic value of fine-needle aspiration biopsy for breast mass: a systematic review and meta-analysis. *BMC Cancer* 2012;12: 41.
17. Abu- Salem OT. Fine needle aspiration biopsy (FNAB) of breast lump: comparison study between pre and post-operative histological diagnosis. *Arch Inst Pasteur Tunis* 2002;79: 59-63.
18. Ibikunle DE, Omotayo JA, Ariyibi OO. Fine needle aspiration cytology of breast lumps with histopathologic correlation in Owo, Ondo State, Nigeria: a five-year review. *Ghana Med J*. 2017; 51 (1):1-5.
19. Lieske B, Ravichandran D, and Wright D. Role of fine-needle aspiration cytology and core biopsy in the preoperative diagnosis of screen-detected breast carcinoma. *Br J Cancer*. 2006; 95: 62–6.
20. Britton PD, Flower CD, Freeman AH, Sinnatambay R, Warren R, Goddard MJ et al. Changing to core biopsy in an NHS breast screening unit. *Clin Radiol* 1997; 52:764-767.
21. Anwar M R, Saeed N, Shahzad M K. Reliability of Fine Needle Aspiration Cytology in Patients with Palpable Breast Lumps. *P J M H S*. 2013; 7(1): 243–6.
22. Saraf S, Khare M, Kalgutkar A. Fine Needle Aspiration Cytology of Breast Lumps- A correlation with histopathology diagnosis.

- Indian Journal of Pathology and Oncology 2016; 3(1);103-6.
23. Shastri R K. Fine Needle Aspiration Cytology Verses Trucut Biopsy In Preoperative Planning Of Surgical Treatment Of Breast Lumps. IOSR-JDMS. 2016; 15(7): 74-80
24. Laws SA, McNamara IR, Cheetham JE, Rainsbury RM. Fine needle aspiration cytology prior to breast ultrasonography does not alter ultrasound diagnostic accuracy or patient management. Breast J 2002; 11: 320-3.
25. Ljung BM, Drejet A, Chianipi N, Jeffery J, Goodson WH, Chew K. Diagnostic accuracy biopsy us determined by physician training in sampling technique. Ce. Cancer 2001; 93: 265-8.